

## References

- Adams, A. M. (1986). The parasite community on the gills of *Fundulus kansae* (Garman) from the South Platte River, Nebraska (USA). *Acta Parasitologica Polonica*, 31, 47-54.
- Adams, A. M. (1984). Infestation of *Fundulus kansae* (Garman) (pisces: Cyprinodontidae) by the copepod *Lernaea cyprinacea* Linnaeus, 1758, in the South Platte River, Nebraska. *American Midland Naturalist*, 112(1), 131-137.
- Adams, A. M. (1985). Parasites on the gills of the plains killifish, *Fundulus kansae*, in the South Platte River, Nebraska. *Transactions of the American Microscopical Society*, 104(3), 278-284.
- Admiraal, D. M., Stansbury, J. S., & Haberman, C. J. (2004). Case study: Particle velocimetry in a model of Lake Ogallala. *Journal of Hydraulic Engineering*, 130(7), 599-607.  
doi:10.1061/(ASCE)0733-9429(2004)130:7(599)
- Alward, R. D., & Joern, A. (1993). Plasticity and overcompensation in grass responses to herbivory. *Oecologia*, 95(3), 358-364. doi:10.1007/BF00320989
- Anderson, J. A., Blazek, K. J., Percival, T. J., & Janovy Jr, J. (1993). The niche of the gill parasite *Dactylogyrus banghami* (monogenea: Dactylogyridae) on *Notropis stramineus* (pisces: Cyprinidae). *The Journal of Parasitology*, 79(3), 435-437.
- Appel, H. M., & Joern, A. (1998). Gut physicochemistry of grassland grasshoppers. *Journal of Insect Physiology*, 44(7-8), 693-700. doi:10.1016/S0022-1910(97)00122-4

Arnett, A. E., & Louda, S. M. (2002). Re-test of *Rhinocyllus conicus* host specificity, and the prediction of ecological risk in biological control. *Biological Conservation*, 106(2), 251-257. doi:10.1016/S0006-3207(01)00251-8

Baack, E. J., Sapir, Y., Chapman, M. A., Burke, J. M., & Rieseberg, L. H. (2008). Selection on domestication traits and quantitative trait loci in crop-wild sunflower hybrids. *Molecular Ecology*, 17(2), 666-677. doi:10.1111/j.1365-294X.2007.03596.x

Ballinger, R. E., & Jones, S. M. (1985). Ecological disturbance in a sandhills prairie: Impact and importance to the lizard community on Arapaho Prairie in western Nebraska. *Prairie Naturalist*, 17(2), 91-100.

Ballinger, R. E., Jones, S. M., & Nietfeldt, J. W. (1990). Patterns of resource use in a lizard community in the Nebraska sandhill prairie. *Prairie Naturalist*, 22(2), 75-86.

Ballinger, R. E., & Watts, K. S. (1995). Path to extinction: Impact of vegetational change on lizard populations on Arapaho Prairie in the Nebraska Sandhills. *American Midland Naturalist*, 134(2), 413-417.

Ballinger, R. E., Droege, D. L., & Jones, S. M. (1981). Reproduction in a Nebraska sandhills population of the northern prairie lizard *Sceloporus undulatus garmani*. *American Midland Naturalist*, 106(1), 157-164.

Ballinger, R. E., Lynch, J. D., & Cole, P. H. (1979). Distribution and natural history of amphibians and reptiles in western Nebraska with ecological notes on the herpetiles of Arapaho Prairie. *Prairie Naturalist*, 11(3), 65-74.

Ballinger, R. E., Nietfeldt, J. W., & Krupa, J. J. (1979). An experimental analysis of the role of the tail in attaining high running speed in *Cnemidophorus sexlineatus* (reptilia: Squamata: Lacertilia). *Herpetologica*, 35(2), 114-116.

Bandi, C., La Rosa, G., Bardin, M. G., Damiani, G., de Carneri, I., & Pozio, E. (1993).

Arbitrarily primed polymerase chain reaction of individual *Trichinella* specimens. *The Journal of Parasitology*, 79(3), 437-440.

Barceló, I., & Faaborg, J. (2012). Expansion of the breeding range of the blue-gray gnatcatcher (*Polioptila caerulea*) into western Nebraska. *The Southwestern Naturalist*, 57(4), 478-479.  
doi:10.1894/0038-4909-57.4.478

Barger, M. A., & Janovy Jr, J. J. (1994). Host specificity of *Rhabdochona canadensis* (nematoda: Rhabdochonidae) in Nebraska. *The Journal of Parasitology*, 80(6), 1032-1035.

Barger, M. A., & Nickol, B. B. (1998). Structure of *Leptorhynchoides thecatus* and *Pomphorhynchus bulbocolli* (acanthocephala) eggs in habitat partitioning and transmission. *The Journal of Parasitology*, 84(3), 534-537.

Barnes, P. W. (1986). Variation in the big bluestem (*Andropogon gerardii*)-sand bluestem (*Andropogon hallii*) complex along a local dune/meadow gradient in the Nebraska Sandhills. *American Journal of Botany*, 73(2), 172-184.

Barnes, M. E., & Durben, D. J. (2010). An evaluation of DVaqua®, a fully-fermented yeast culture, during long-term hatchery rearing of McConaughy strain rainbow trout. *Aquaculture Nutrition*, 16(3), 299-304. doi:10.1111/j.1365-2095.2009.00665.x

Barnes, P. W., Harrison, A. T., & Heinisch, S. P. (1984). Vegetation patterns in relation to topography and edaphic variation Nebraska sandhills prairie. *Prairie Naturalist*, 16(4), 145-158.

Barnes, P. W. (1985). Adaptation to water stress in the big bluestem-sand bluestem complex. *Ecology*, 66(6), 1908-1920. doi:10.2307/2937387

Barnes, P. W., & Harrison, A. T. (1982). Species distribution and community organization in a Nebraska sandhills mixed prairie as influenced by plant/soil-water relationships. *Oecologia*, 52(2), 192-201. doi:10.1007/BF00363836

Baxa, M. D. (1981). Anodonta species of Lake McConaughy, western Nebraska. *The Nautilus*, 95(4), 180-183.

Behmer, S. T., & Joern, A. (1993). Diet choice by a grass-feeding grasshopper based on the need for a limiting nutrient. *Functional Ecology*, 7(5), 522-527.

Behmer, S. T., & Joern, A. (1993). Dietary selection by the generalist grasshopper, *Phoetaliotes nebrascensis* (orthoptera: Acrididae) based on the need for phenylalanine. *Functional Ecology*, 7, 522-527.

Behmer, S. T., & Joern, A. (1994). The influence of proline on diet selection: Sex-specific feeding preferences by the grasshoppers *Ageneotettix deorum* and *Phoetaliotes nebrascensis* (orthoptera: Acrididae). *Oecologia*, 98(1), 76-82. doi:10.1007/BF00326093

Behmer, S. T., & Joern, A. (2008). Coexisting generalist herbivores occupy unique nutritional feeding niches. *Proceedings of the National Academy of Sciences*, 105(6), 1977-1982. doi:10.1073/pnas.0711870105

Belovsky, G. E., Botkin, D. B., Crowl, T. A., Cummins, K. W., Franklin, J. F., Hunter Jr, M., L., . . Margules, C. R. (2004). Ten suggestions to strengthen the science of ecology. *Bioscience*, 54(4), 345-351. doi:10.1641/0006-3568(2004)054[0345:TSTSTS]2.0.CO;2

Benedict, R. A., Genoways, H. H., & Freeman, P. W. (2000). Shifting distributional patterns of mammals in Nebraska. *Transactions of the Nebraska Academy of Sciences*, 26, 55-84.

Bergin, T. M. (1992). Cost of nest reuse by western kingbirds. *Prairie Naturalist*, 241, 39-40.

Bergin, T. M. (1992). Habitat selection by the western kingbird in western Nebraska: A hierarchical analysis. *The Condor*, 94(4), 903-911.

Bergin, T. M. (1997). Nest reuse by western kingbirds. *The Wilson Bulletin*, 109(4), 735-737.

Bergin, T. M., & Paseka, J. (1997). Birds of the Lake Ogallala State Recreation Area. *Nebraska Bird Review*, 65(2), 59-67.

Bevill, R. L., & Louda, S. M. (1999). Comparisons of related rare and common species in the study of plant rarity. *Conservation Biology*, 13(3), 493-498. doi:10.1046/j.1523-1739.1999.97369.x

Bevill, R. L., Louda, S. M., & Stanforth, L. M. (1999). Protection from natural enemies in managing rare plant species. *Conservation Biology*, 13(6), 1323-1331. doi:10.1046/j.1523-1739.1999.98450.x

Blackman, B. K., Rasmussen, D. A., Strasburg, J. L., Raduski, A. R., Burke, J. M., Knapp, S. J., Rieseberg, L. H. (2011). Contributions of flowering time genes to sunflower domestication and improvement. *Genetics*, 187(1), 271-287. doi:10.1534/genetics.110.121327

Bock, C. E., & Scharf, W. C. (1994). A nesting population of cassin's sparrows in the sandhills of Nebraska. *Journal of Field Ornithology*, 65(4), 472-475.

Bolek, M. G., Brotan, K. K., Rudolph, R. E., & Janovy Jr, J. J. (2007). *Bufo woodhousii* (woodhouse's toad). cannibalism. *Herpetological Review*, 38(3), 319.

Bolek, M. G., & Janovy Jr, J. (2007). Evolutionary avenues for, and constraints on, the transmission of frog lung flukes (*Haematoloechus* spp.) in dragonfly second intermediate hosts. *The Journal of Parasitology*, 93(3), 593-607. doi:10.1645/GE-1011R.1

Bolek, M. G., & Janovy Jr, J. (2008). Alternative life cycle strategies of *Megalodiscus temperatus* in tadpoles and metamorphosed anurans. *Parasite*, 15(3), 396-401. doi:10.1051/parasite/2008153396

Bolek, M. G., & Janovy Jr, J. (2005). New host and distribution records for the amphibian leech *Desserobdella picta* (rhynchobdellida: Glossiphoniidae) from Nebraska and Wisconsin. *Journal of Freshwater Ecology*, 20(1), 187-189. doi:10.1080/02705060.2005.9664951

Bolek, M. G., & Janovy Jr, J. (2007). Small frogs get their worms first: The role of nonodonate arthropods in the recruitment of *Haematoloechus coloradensis* and *Haematoloechus complexus* in newly metamorphosed northern leopard frogs, *Rana pipiens*, and woodhouse's toads, *Bufo woodhousii*. *Journal of Parasitology*, 93(2), 300-312. doi:10.1645/GE-1010R.1

Bolek, M. G., Janovy Jr, J., & Irizarry-Rovira, A. (2003). Observations on the life history and descriptions of coccidia (apicomplexa) from the western chorus frog, *Pseudacris triseriata triseriata*, from eastern Nebraska. *The Journal of Parasitology*, 89(3), 522-528. doi:10.1645/0022-3395(2003)089[0522:OOTLHA]2.0.CO;2

Bolek, M. G., Rogers, E., Szmygiel, C., Shannon, R. P., Doerfert-Schrader, W., Schmidt-Rhaesa, A., & Hanelt, B. (2013). Survival of larval and cyst stages of gordiids (nematomorpha) after exposure to freezing. *The Journal of Parasitology*, 99(3), 397-402. doi:10.1645/12-62.1

Bolek, M. G., & Schmidt-Rhaesa, A. (2010). Redescription of the African *Chordodes albibarbatus* Montgomery 1898, and description of *Chordodes janovyi* n. sp.(gordiida, nematomorpha) and its non-adult stages from Cameroon, Africa. *Zootaxa*, 2631, 36-50.

Bolek, M. G., Snyder, S. D., & Janovy Jr, J. (2009). Alternative life cycle strategies and colonization of young anurans by *Gorgoderina attenuata* in Nebraska. *Journal of Parasitology*, 95(3), 604-616. doi:10.1645/GE-1813.1

Bolek, M. G., Snyder, S. D., & Janovy Jr, J. (2009). Redescription of the frog bladder fluke *Gorgoderina attenuata* from the northern leopard frog, *Rana pipiens*. *Journal of Parasitology*, 95(3), 665-668. doi:10.1645/GE-1953.1

Bolek, M. G., Szmygiel, C., Kubat, A., Schmidt-Rhaesa, A., & Hanelt, B. (2013). Novel techniques for biodiversity studies of gordiids and description of a new species of chordodes (gordiida, nematomorpha) from Kenya, Africa. *Zootaxa*, 3717(1), 023-038. doi:10.11646/zootaxa.3717.1.2

Bolek, M. G., Tracy, H. R., & Janovy Jr, J. (2010). The role of damselflies (odonata: Zygoptera) as paratenic hosts in the transmission of *Halipegus eccentricus* (digenea: Hemiuridae) to anurans. *Journal of Parasitology*, 96(4), 724-735. doi:10.1645/GE-2365.1

Bomberger Brown, M., & Brown, C. R. (2011). Intense natural selection on morphology of cliff swallows (*Petrochelidon pyrrhonota*) a decade later: Did the population move between adaptive peaks? *The Auk*, 128(1), 69-77. doi:10.1525/auk.2011.10219

Bomberger, M. L. (1984). Quantitative assessment of the nesting habitat of Wilson's phalarope. *The Wilson Bulletin*, 96(1), 126-128.

Bomberger-Brown, M.,L., Shields, S. L., Harrsion, A. T., & Keeler, K. H. (1983). Comparison of old field succession on a tallgrass prairie and a Nebraska sandhills prairie. *Prairie Naturalist*, 13(1), 9-15.

Bradley, K. L., Damschen, E. I., Young, L. M., Kuefler, D., Went, S., Wray, G., Louda, S. M. (2003). Spatial heterogeneity, not visitation bias, dominates variation in herbivory. *Ecology*, 84(8), 2214-2221.

Branson, D. H., Joern, A., & Sword, G. A. (2006). Sustainable management of insect herbivores in grassland ecosystems: New perspectives in grasshopper control. *Bioscience*, 56(9), 743-755. doi:10.1641/0006-3568(2006)56[743:SMOIHI]2.0.CO;2

Brown, C. R. (1988). Enhanced foraging efficiency through information centers: A benefit of coloniality in cliff swallows. *Ecology*, 69(3), 602-613. doi:10.2307/1941009

Brown, C. R., & Brown, M. B. (1988). A new form of reproductive parasitism in cliff swallows. *Nature*, 331(6151), 66-68. doi:10.1038/331066a0

Brown, C. R., & Brown, M. B. (1989). Behavioural dynamics of intraspecific brood parasitism in colonial cliff swallows. *Animal Behaviour*, 37(5), 777-796. doi:10.1016/0003-3472(89)90063-8

Brown, C. R., & Brown, M. B. (1992). Ectoparasitism as a cause of natal dispersal in cliff swallows. *Ecology*, 73(5), 1718-1723. doi:10.2307/1940023

Brown, C. R., & Brown, M. B. (1998). Intense natural selection on body size and wing and tail asymmetry in cliff swallows during severe weather. *Evolution*, 52(5), 1461-1475.

Brown, C. R., & Brown, M. B. (1999). Fitness components associated with clutch size in cliff swallows. *The Auk*, 116(2), 467-486.

Brown, C. R., & Brown, M. B. (2004). Mark-recapture and behavioral ecology: A case study of cliff swallows. *Animal Biodiversity and Conservation*, 27(1), 23-34.

Brown, C. R., Brown, M. B., & Ives, A. R. (1992). Nest placement relative to food and its influence on the evolution of avian coloniality. *The American Naturalist*, 139(1), 205-217.

Brown, C. R., Brown, M. B., Moore, A. T., & Komar, N. (2007). Bird movement predicts buggy creek virus infection in insect vectors. *Vector Borne and Zoonotic Diseases*, 7(3), 304-314. doi:10.1089/vbz.2006.0646

Brown, C. R., Brown, M. B., & Roche, E. A. (2013). Fluctuating viability selection on morphology of cliff swallows is driven by climate. *Journal of Evolutionary Biology*, 26(5), 1129-1142. doi:10.1111/jeb.12130

Brown, C. R., & Sherman, L. C. (1989). Variation in the appearance of swallow eggs and the detection of intraspecific brood parasitism. *The Condor*, 91(3), 620-627.

Brown, C. R. (1984). Laying eggs in a neighbor's nest: Benefit and cost of colonial nesting in swallows. *Science*, 224(4648), 518-519. doi:10.1126/science.224.4648.518

Brown, C. R. (1985). Vocalizations of barn and cliff swallows. *The Southwestern Naturalist*, 30(3), 325-333.

Brown, C. R. (1986). Cliff swallow colonies as information centers. *Science*, 234(4772), 83-85. doi:10.1126/science.234.4772.83

Brown, C. R. (1988). Social foraging in cliff swallows: Local enhancement, risk sensitivity, competition and the avoidance of predators. *Animal Behaviour*, 36(3), 780-792. doi:10.1016/S0003-3472(88)80161-1

Brown, C. R. (1990). Social foraging in cliff swallows revisited. *Animal Behaviour*, 39(6), 1216-1218.

Brown, C. R. (2003). Birds of Nebraska: Their distribution and temporal occurrence. *The Condor*, 105(3), 609-610. doi:10.1650/7349

Brown, C. R., Brazeal, K. R., Strickler, S. A., & Brown, M. B. (2006). Feather mites are positively associated with daily survival in cliff swallows. *Canadian Journal of Zoology*, 84(9), 1307-1314. doi:10.1139/z06-110

Brown, C. R., & Brown, M. B. (1992). First record of the cave swallow for Nebraska. *Nebraska Bird Review*, 60(1), 36-40.

Brown, C. R., & Brown, M. B. (2013). Where has all the road kill gone? *Current Biology*, 23(6), R233-R234. doi:10.1016/j.cub.2013.02.023

Brown, C. R., & Brown, M. B. (1986). Ectoparasitism as a cost of coloniality in cliff swallows (*Hirundo pyrrhonota*). *Ecology*, 67(5), 1206-1218.

Brown, C. R., & Brown, M. B. (1987). Group-living in cliff swallows as an advantage in avoiding predators. *Behavioral Ecology and Sociobiology*, 21(2), 97-107.

Brown, C. R., & Brown, M. B. (1988). The costs and benefits of egg destruction by conspecifics in colonial cliff swallows. *The Auk*, 105(4), 737-748.

Brown, C. R., & Brown, M. B. (1988). Genetic evidence of multiple parentage in broods of cliff swallows. *Behavioral Ecology and Sociobiology*, 23(6), 379-387.

Brown, C. R., & Brown, M. B. (1990). The great egg scramble. *Natural History*, (2), 34-41.

Brown, C. R., & Brown, M. B. (1991). Selection of high-quality host nests by parasitic cliff swallows. *Animal Behaviour*, 41(3), 457-465.

Brown, C. R., & Brown, M. B. (1998). Fitness components associated with alternative reproductive tactics in cliff swallows. *Behavioral Ecology*, 9(2), 158-171.  
doi:10.1093/beheco/9.2.158

Brown, C. R., & Brown, M. B. (1999). Fitness components associated with laying date in the cliff swallow. *Condor*, 101(2), 230-245.

Brown, C. R., & Brown, M. B. (1999). Natural selection on tail and bill morphology in barn swallows *Hirundo rustica* during severe weather. *Ibis*, 141(4), 652-659. doi:10.1111/j.1474-919X.1999.tb07373.x

Brown, C. R., & Brown, M. B. (2000). Heritable basis for choice of group size in a colonial bird. *Proceedings of the National Academy of Sciences*, 97(26), 14825-14830.  
doi:10.1073/pnas.97.26.14825

Brown, C. R., & Brown, M. B. (2000). Nest spacing in relation to settlement time in colonial cliff swallows. *Animal Behaviour*, 59(1), 47-55. doi:10.1006/anbe.1999.1277

Brown, C. R., & Brown, M. B. (2000). Weather-mediated natural selection on arrival time in cliff swallows (*Petrochelidon pyrrhonota*). *Behavioral Ecology and Sociobiology*, 47(5), 339-345. doi:10.1007/s002650050674

Brown, C. R., & Brown, M. B. (2001). Egg hatchability increases with colony size in cliff swallows. *Journal of Field Ornithology*, 72(1), 113-123. doi:10.1648/0273-8570-72.1.113

Brown, C. R., & Brown, M. B. (2002). Does intercolony competition for food affect colony choice in cliff swallows? *The Condor*, 104(1), 117-128. doi:10.1650/0010-5422(2002)104[0117:DICFFA]2.0.CO;2

Brown, C. R., & Brown, M. B. (2002). Ectoparasites cause increased bilateral asymmetry of naturally selected traits in a colonial bird. *Journal of Evolutionary Biology*, 15(6), 1067-1075. doi:10.1046/j.1420-9101.2002.00474.x

Brown, C. R., & Brown, M. B. (2002). Spleen volume varies with colony size and parasite load in a colonial bird. *Proceedings of the Royal Society of London.Series B: Biological Sciences*, 269(1498), 1367-1373. doi:10.1098/rspb.2002.2036

Brown, C. R., & Brown, M. B. (2003). Testis size increases with colony size in cliff swallows. *Behavioral Ecology*, 14(4), 569-575. doi:10.1093/beheco/arg030

Brown, C. R., & Brown, M. B. (2004). Empirical measurement of parasite transmission between groups in a colonial bird. *Ecology*, 85(6), 1619-1626. doi:10.1890/03-0206

Brown, C. R., & Brown, M. B. (2004). Group size and ectoparasitism affect daily survival probability in a colonial bird. *Behavioral Ecology and Sociobiology*, 56(5), 498-511. doi:10.1007/s00265-004-0813-6

Brown, C. R., & Brown, M. B. (2005). Between-group transmission dynamics of the swallow bug, *Oeciacus vicarius*. *Journal of Vector Ecology*, 30(1), 137-143.

Brown, C. R., Brown, M. B., & Brazeal, K. R. (2008). Familiarity with breeding habitat improves daily survival in colonial cliff swallows. *Animal Behaviour*, 76(4), 1201-1210. doi:10.1016/j.anbehav.2008.03.028

Brown, C. R., Brown, M. B., & Danchin, E. (2000). Breeding habitat selection in cliff swallows: The effect of conspecific reproductive success on colony choice. *Journal of Animal Ecology*, 69(1), 133-142. doi:10.1046/j.1365-2656.2000.00382.x

Brown, C. R., Brown, M. B., Johnsgard, P. A., Kren, J., & Scharf, W. C. (1996). Birds of the Cedar Point Biological Station area, Keith and Garden counties, Nebraska: Seasonal occurrence and breeding data. *Transactions of the Nebraska Academy of Sciences*, 23, 91-108.

Brown, C. R., Brown, M. B., Padhi, A., Foster, J. E., Moore, A. T., Pfeffer, M., & Komar, N. (2008). Host and vector movement affects genetic diversity and spatial structure of buggy creek virus (togaviridae). *Molecular Ecology*, 17(9), 2164-2173. doi:10.1111/j.1365-294X.2008.03747.x

Brown, C. R., Brown, M. B., & Rannala, B. (1995). Ectoparasites reduce long-term survival of their avian host. *Proceedings of the Royal Society of London.Series B: Biological Sciences*, 262(1365), 313-319. doi:10.1098/rspb.1995.0211

Brown, C. R., Brown, M. B., Raouf, S. A., Smith, L. C., & Wingfield, J. C. (2005). Effects of endogenous steroid hormone levels on annual survival in cliff swallows. *Ecology*, 86(4), 1034-1046. doi:10.1890/04-0740

Brown, C. R., Brown, M. B., Raouf, S. A., Smith, L. C., & Wingfield, J. C. (2005). Steroid hormone levels are related to choice of colony size in cliff swallows. *Ecology*, 86(11), 2904-2915. doi:10.1890/05-0246

Brown, C. R., Brown, M. B., & Roche, E. A. (2013). Spatial and temporal unpredictability of colony size in cliff swallows across 30 years. *Ecological Monographs*, 83, 511-530. doi:10.1890/12-2001.1

Brown, C. R., Brown, M. B., & Shaffer, M. L. (1991). Food-sharing signals among socially foraging cliff swallows. *Animal Behaviour*, 42(4), 551-564.

Brown, C. R., & Hoogland, J. L. (1986). Risk in mobbing for solitary and colonial swallows. *Animal Behaviour*, 34(5), 1319-1323.

Brown, C. R., Komar, N., Quick, S. B., Sethi, R. A., Panella, N. A., Brown, M. B., & Pfeffer, M. (2001). Arbovirus infection increases with group size. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 268(1478), 1833-1840. doi:10.1098/rspb.2001.1749

Brown, C. R., Moore, A. T., & O'Brien, V.,A. (2012). Prevalence of buggy creek virus (togaviridae: *Alphavirus*) in insect vectors increases over time in the presence of an invasive avian host. *Vector-Borne and Zoonotic Diseases*, 12(1), 34-41. doi:10.1089/vbz.2011.0677

Brown, C. R., Moore, A. T., O'Brien, V.,A., Padhi, A., Knutie, S. A., Young, G. R., & Komar, N. (2010). Natural infection of vertebrate hosts by different lineages of buggy creek virus (family togaviridae, genus *Alphavirus*). *Archives of Virology*, 155(5), 745-749. doi:10.1007/s00705-010-0638-8

Brown, C. R., Moore, A. T., Young, G. R., & Komar, N. (2010). Persistence of buggy creek virus (togaviridae, *Alphavirus*) for two years in unfed swallow bugs (hemiptera: Cimicidae: *Oeciacus vicarius*). *Journal of Medical Entomology*, 47(3), 436-441.

Brown, C. R., Moore, A. T., Young, G. R., Padhi, A., & Komar, N. (2009). Isolation of buggy creek virus (togaviridae: *Alphavirus*) from field-collected eggs of *Oeciacus vicarius* (hemiptera: Cimicidae). *Journal of Medical Entomology*, 46(2), 375-379.

Brown, C. R., Padhi, A., Moore, A. T., Brown, M. B., Foster, J. E., Pfeffer, M., . . . Komar, N. (2009). Ecological divergence of two sympatric lineages of buggy creek virus, an arbovirus associated with birds. *Ecology*, 90(11), 3168-3179. doi:10.1890/08-1731.1

Brown, C. R., & Rannala, B. (1995). Colony choice in birds: Models based on temporally invariant site quality. *Behavioral Ecology and Sociobiology*, 36(4), 221-228.

Brown, C. R., Sas, C. M., & Brown, M. B. (2002). Colony choice in cliff swallows: Effects of heterogeneity in foraging habitat. *The Auk*, 119(2), 446-460. doi:10.1642/0004-8038(2002)119[0446:CCICSE]2.0.CO;2

Brown, C. R., & Sethi, R. A. (2002). Mosquito abundance is correlated with cliff swallow (*Petrochelidon pyrrhonota*) colony size. *Journal of Medical Entomology*, 39(1), 115-120. doi:10.1603/0022-2585-39.1.115

Brown, C. R., Strickler, S. A., Moore, A. T., Knutie, S. A., Padhi, A., Brown, M. B., . . . Komar, N. (2010). Winter ecology of buggy creek virus (togaviridae, *Alphavirus*) in the central great plains. *Vector-Borne and Zoonotic Diseases*, 10(4), 355-363. doi:10.1089/vbz.2009.0031

Brown, C. R., Stutchbury, B. J., & Walsh, P. D. (1990). Choice of colony size in birds. *Trends in Ecology & Evolution*, 5(12), 398-403.

Brown, C. R., & Brown, M. B. (1995). Cliff swallow: *Hirundo pyrrhonota*. In A. Poole, & F. Gill (Eds.), *Birds of North America* (pp. 32). Philadelphia ad Washington D.C.: Academy of Natural Sciences and American Ornithologists' Union.

Brown, M. B., & Brown, C. R. (2009). Blood sampling reduces annual survival in cliff swallows (*Petrochelidon pyrrhonota*). *The Auk*, 126(4), 853-861. doi:10.1525/auk.2009.09048

Bunker, B. E., Janovy Jr, J., Tracey, E., Barnes, A., Duba, A., Shuman, M., & Logan, J. D. (2013). Macroparasite population dynamics among geographical localities and host life cycle stages: Eugregarines in *Ischnura verticalis*. *The Journal of Parasitology*, 99(3), 403-409. doi:10.1645/GE-3137.1

Burger, J. C., & Louda, S. M. (1994). Indirect versus direct effect of grasses on growth of a cactus (*Opuntia fragilis*): Insect herbivory versus competition. *Oecologia*, 99(1-2), 79-87.

Burger, J. C., & Louda, S. M. (1995). Interaction of diffuse competition and insect herbivory in limiting brittle prickly pear cactus, *Opuntia fragilis* (cactaceae). *American Journal of Botany*, 82(12), 1558-1566.

Carter, M. R. (1989). The biology and ecology of the tiger beetles (coleoptera: Cincindelidae) of Nebraska. *Transactions of the Nebraska Academy of Sciences*, 17, 1-18.

Clopton, R. E. (1991). A review of the scaritiniid beetles (coleoptera: Carabidae: Scaritini). *Transactions of the Nebraska Academy of Sciences and Affiliated Societies*, 18, 53-65.

Clopton, R. E. (1999). Revision of the genus *Xiphoccephalus* and description of *Xiphoccephalus ellisi* n. sp.(apicomplexa: Eugregarinida: Stylocephalidae) from *Eleodes opacus* (coleoptera: Tenebrionidae) in the western Nebraska sandhills. *The Journal of Parasitology*, 85(1), 84-89.

Clopton, R. E. (2000). *Stylocephalus occidentalis* n. sp.(apicomplexa: Eugregarinida: Stylocephalidae) from *Trimytilis pruinosa* (coleoptera: Tenebrionidae) in the Nebraska sandhills. *The Journal of Parasitology*, 86(3), 560-565. doi:10.1645/0022-3395(2000)086[0560:SONSAE]2.0.CO;2

Clopton, R. E. (2006). Two new species of *Xiphoccephalus* in *Eleodes tricostata* and *Eleodes fusiformis* (coleoptera: Tenebrionidae: Eleodini) from the sandhills of western Nebraska. *Journal of Parasitology*, 92(3), 569-577. doi:10.1645/GE-742R.1

Clopton, R. E., Janovy Jr, J., & Percival, T. J. (1992). Host stadium specificity in the gregarine assemblage parasitizing *Tenebrio molitor*. *The Journal of Parasitology*, 78(2), 334.

Clopton, R. E., & Janovy Jr, J. J. (1993). Developmental niche structure in the gregarine assemblage parasitizing *Tenebrio molitor*. *Journal of Parasitology*, 79(5), 701-709.

Clopton, R. E., & Nolte, C. M. (2002). *Clitellocephalus americanus* n. gen., n. sp.(apicomplexa: Eugregarinida: Gregarinidae) from *Cratacanthus dubius* (coleoptera: Carabidae: Harpalinae) in the Nebraska sandhills and *Clitellocephalus ophoni* n. comb.(apicomplexa: Eugregarinida: Gregarinidae) from *Ophonus pubescens* (coleoptera: Carabidae: Harpalinae) in Sète, France. *The Journal of Parasitology*, 88(4), 750-757. doi:10.1645/0022-3395(2002)088[0750:CANGNS]2.0.CO;2

Clopton, R. E., Percival, T. J., & Janovy Jr, J. (1993). *Nubenocephalus nebraskensis* n. gen., n. sp.(apicomplexa: Actinocephalidae) from adults of *Argia bipunctulata* (odonata: Zygoptera). *The Journal of Parasitology*, 79(4), 533-537.

Clopton, R. E., Percival, T. J., & Janovy, J. (1992). *Gregarina coronata* N. sp. (apicomplexa: Eugregarinida) described from adults of the southern corn rootworm, *Diabrotica undecimpunctata howardi* (coleoptera: Chrysomelidae). *Journal of Eukaryotic Microbiology*, 39(3), 417-420. doi:10.1111/j.1550-7408.1992.tb01474.x

Clopton, R. E., Cook, T. J., & Cielocha, J. J. (2010). *Nubenocephalus nickoli* n. sp. and *Nubenocephalus xunantunichensis* n. sp.(apicomplexa: Eugregarinida: Actinocephalidae) parasitizing damselflies (odonata: Zygoptera) in Belize, Central America. *Comparative Parasitology*, 77(2), 125-136. doi:10.1654/4438.1

Clopton, R. E., Percival, T. J., & Janovy, J. (1991). *Gregarina niphandrodes* n. sp.(apicomplexa: Eugregarinorida) from adult *Tenebrio molitor* (L.) with oocyst descriptions of other gregarine parasites of the yellow mealworm. *Journal of Eukaryotic Microbiology*, 38(5), 472-479.

Coelho, J. R., Hastings, J. M., Holliday, C. W., Flure, G., & Barnes, K. (2008). Sexual dimorphism of the femora, tibiae, and hind tibial spurs in the eastern cicada killer, *Sphecius speciosus* Drury (hymenoptera: Crabronidae) in the United States. *Entomological News*, 119(1), 11-18. doi:10.3157/0013-872X(2008)119[11:SDOTFT]2.0.CO;2

Collins, M. R., & Janovy Jr, J. (2003). Host specificity among ancyrocephalinae (monogenoidea) of Nebraska sunfish. *Journal of Parasitology*, 89(1), 80-83. doi:10.1645/0022-3395(2003)089[0080:HSAAMO]2.0.CO;2

Current, W. L., & Janovy Jr, J. (1977). Sporogenesis in *Henneguya exilis* infecting the channel catfish: An ultrastructural study. *Protistologica*, 13, 157-167.

Current, W. L., & Janovy Jr, J. (1976). Ultrastructure of interlamellar *Henneguya exilis* in the channel catfish. *The Journal of Parasitology*, 62(6), 975-981.

Current, W. L., & Janovy Jr, J. (1978). Comparative study of ultrastructure of interlamellar and intralamellar types of *Henneguya exilis* kudo from channel catfish. *The Journal of Protozoology*, 25(1), 56-65.

Current, W. L., Janovy Jr, J., & Knight, S. A. (1979). *Myxosoma funduli* kudo (myxosporida) in *Fundulus kansae*: Ultrastructure of the plasmodium wall and of sporogenesis. *The Journal of Protozoology*, 26(4), 574-583.

Danner, B. J., & Joern, A. (2003). Resource-mediated impact of spider predation risk on performance in the grasshopper *Ageneotettix deorum* (orthoptera: Acrididae). *Oecologia*, 137(3), 352-359. doi:10.1007/s00442-003-1362-9

Danner, B. J., & Joern, A. (2003). Stage-specific behavioral responses of *Ageneotettix deorum* (orthoptera: Acrididae) in the presence of lycosid spider predators. *Journal of Insect Behavior*, 16(4), 453-464. doi:10.1023/A:1027337405495

Danner, B., & Joern, A. (2004). Development, growth, and egg production of *Ageneotettix deorum* (orthoptera: Acrididae) in response to spider predation risk and elevated resource quality. *Ecological Entomology*, 29(1), 1-11. doi:10.1111/j.0307-6946.2004.00571.x

David Logan, J., Joern, A., & Wolessensky, W. (2002). Location, time, and temperature dependence of digestion in simple animal tracts. *Journal of Theoretical Biology*, 216(1), 5-18.

Dechaine, J. M., Burger, J. C., & Burke, J. M. (2010). Ecological patterns and genetic analysis of post-dispersal seed predation in sunflower (*Helianthus annuus*) crop-wild hybrids. *Molecular Ecology*, 19(16), 3477-3488. doi:10.1111/j.1365-294X.2010.04740.x

Dechaine, J. M., Burger, J. C., Chapman, M. A., Seiler, G. J., Brunick, R., Knapp, S. J., & Burke, J. M. (2009). Fitness effects and genetic architecture of plant-herbivore interactions in sunflower crop-wild hybrids. *New Phytologist*, 184(4), 828-841. doi:10.1111/j.1469-8137.2009.02964.x

Decker, K. L., & Pilson, D. (2000). Biased sex ratios in the dioecious annual *Croton texensis* (euphorbiaceae) are not due to environmental sex determination. [Environmental sex determination as a possible cause of biased sex ratios in *Croton texensis*] *American Journal of Botany*, 87(2), 221-229.

DeNicola, M., & Hoagland, K. D. (1996). Effects of solar spectral irradiance (visible to UV) on a prairie stream epilithic community. *Journal of the North American Benthological Society*, 15(2), 155-169.

DeNicola, M., Hoagland, K. D., & Roemer, S. C. (1992). Influences of canopy cover on spectral irradiance and periphyton assemblages in a prairie stream. *Journal of the North American Benthological Society*, 11(4), 391-404.

Diffendal, R. F. (1982). Regional implications of the geology of the Ogallala group (upper tertiary) of southwestern Morrill County, Nebraska, and adjacent areas. *Geological Society of America Bulletin*, 93(10), 964-976.

Dinsmore, S. J., & Silcock, W. R. (1998). Parasitic jaeger at Lake McConaughy. *Nebraska Bird Review*, 66(4), 114-115.

Droge, D. L., Jones, S. M., & Ballinger, R. E. (1982). Reproduction of *Holbrookia maculata* in western Nebraska. *Copeia*, 1982(2), 356-362.

Durden, L. A., & Richardson, D. J. (2013). Ectoparasites of the virginia opossum (*Didelphis virginiana*), raccoon (*Procyon lotor*), and striped skunk (*Mephitis mephitis*) from Keith County, Nebraska. *Transactions of the Nebraska Academy of Sciences*, 33, 21-24.

Ewel, J. J., O'Dowd, D., Bergelson, J., Daehler, C. C., D'Antonio, C., Gomez, L., . . . Hopper, K. R. (1999). Deliberate introductions of species: Research needs. Benefits can be reaped, but risks are high. *Bioscience*, 49(8), 619-630.

Fassbinder-Orth, C., Barak, V. A., & Brown, C. R. (2013). Immune responses of a native and an invasive bird to buggy creek virus (*togaviridae: Alphavirus*) and its arthropod vector, the swallow bug (*Oeciacus vicarius*). *Plos One*, 8(2), e58045.

doi:10.1371/journal.pone.0058045

- Ferdig, M. T., McDowell, M. A., & Janovy Jr, J. J. (1991). *Salsuginus yutanensis* n. sp. (monogenea: Ancyrocephalidae) from *Fundulus sciadicus* in clear creek of eastern Nebraska. *Journal of Parasitology*, 77(1), 58-61.
- Fertig, M. T., McDowell, M. A., Janovy Jr, J., & Clopton, R. E. (1993). Patterns of morphological variation of *Salsuginus yutanensis* (monogenea: Ancyrocephalidae) over space and time. *The Journal of Parasitology*, 79(5), 744-750.
- Fuller, R., & Joern, A. (1996). Grasshopper susceptibility to predation in response to vegetation cover and patch area. *Journal of Orthoptera Research*, 5, 175-183.
- Gaines, S. B. (1991). Body-size and wing-length variation among selected grasshoppers (orthoptera: Acrididae) from Nebraska's sandhills grasslands. *Transactions of the Nebraska Academy of Sciences*, 18, 67-72.
- González, J., Matthews, R. W., & Matthews, J. R. (2005). Comparative development and competitive ability of *Dibrachys pelos* (hymenoptera: Pteromalidae) on various potential hosts. *Florida Entomologist*, 88(1), 49-54. doi:10.1653/0015-4040(2005)088[0049:CDACAO]2.0.CO;2
- Guretzky, J. A., & Louda, S. M. (1997). Evidence for natural biological control: Insects decrease survival and growth of a native thistle. *Ecological Applications*, 7(4), 1330-1340. doi:10.1890/1051-0761(1997)007[1330:EFNBCI]2.0.CO;2
- Hanelt, B., & Janovy Jr, J. (2000). New host and distribution record of *Gordius difficilis* (nematomorpha: Gordioidea) from a vivid metallic ground beetle, *Chlaenius prasinus*

(coleoptera: Carabidae) from western Nebraska, USA. *Journal of the Helminthological Society of Washington*, 67(1), 107-108.

Hanelt, B., & Janovy Jr, J. (2004). Untying a gordian knot: The domestication and laboratory maintenance of a gordian worm, *Paragordius varius* (nematomorpha: Gordiida). *Journal of Natural History*, 38(8), 939-950. doi:10.1080/0022293021000058718

Hanelt, B., & Janovy JR, ,J. (1999). The life cycle of a horsehair worm, *Gordius robustus* (nematomorpha: Gordioidea). *The Journal of Parasitology*, 85(1), 139-141.

Hanelt, B., Bolek, M. G., & Schmidt-Rhaesa, A. (2012). Going solo: Discovery of the first parthenogenetic gordiid (nematomorpha: Gordiida). *PloS One*, 7(4), e34472. doi:10.1371/journal.pone.0034472

Hanelt, B., Grother, L. E., & Janovy Jr, J. (2001). Physid snails as sentinels of freshwater nematomorphs. *Journal of Parasitology*, 87(5), 1049-1053. doi:10.1645/0022-3395(2001)087[1049:PSASOF]2.0.CO;2

Hanelt, B., & Janovy Jr, J. (2004). Life cycle and paratenesis of American gordiids (nematomorpha: Gordiida). *Journal of Parasitology*, 90(2), 240-244. doi:10.1645/GE-78R

Hanelt, B., & Janovy, J. (2003). Spanning the gap: Experimental determination of paratenic host specificity of horsehair worms (nematomorpha: Gordiida). *Invertebrate Biology*, 122(1), 12-18. doi:10.1111/j.1744-7410.2003.tb00068.x

HANELT, B., & JANOVY, J. J. R. (2002). Morphometric analysis of nonadult characters of common species of American gordiids (nematomorpha: Gordioidea). *The Journal of Parasitology*, 88(3), 557-562. doi:10.1645/0022-3395(2002)088[0557:MAONCO]2.0.CO;2

HARDIN, E. L., & JANOVY JR, J. (1988). Population dynamics of *Distoichometra budonis* (cestoda: Nematotaeniidae) in *Bufo woodhousii*. *The Journal of Parasitology*, 74(3), 360-365.

Hart, C. J., & Ivie, M. A. (2013). Synonymy of *Megacyllene comanchei* rice and morris and *Megacyllene angulifera* (casey)(coleoptera: Cerambycidae). *The Coleopterists Bulletin*, 67(2), 89-93. doi:10.1649/0010-065X-67.2.89

Haynes, J. L., & Weekly, J. M. (1992). New distributional records for three fishes in Lincoln County, Nebraska. *Transactions of the Nebraska Academy of Sciences*, 19, 31-32.

Heidorn, T., & Joern, A. (1987). Feeding preference and spatial distribution of grasshoppers (Acrididae) in response to nitrogen fertilization of *Calamovilfa longifolia*. *Functional Ecology*, 1(4), 369-375.

Heidorn, T., & Joern, A. (1984). Differential herbivory on C3 versus C4 grasses by the grasshopper *Ageneotettix deorum* (orthoptera: Acrididae). *Oecologia*, 65(1), 19-25.

Helt, J., Janovy Jr, J., & Ubelaker, J. (2003). *Phyllostomum funduli* n. sp.(trematoda: Gorgoderidae) from *Fundulus sciadicus* cope from Cedar Creek in western Nebraska. *Journal of Parasitology*, 89(2), 346-350. doi:10.1645/0022-3395(2003)089[0346:PFNSTG]2.0.CO;2

Hiskey, R., Holland, R. S., & Hergenrader, G. L. (1981). The occurrence of *Moina hutchinsonii* Brehm (Cladocera) in a Nebraska sandhills lake. *The Southwestern Naturalist*, 26(2), 217-218.

Hoagland, K. D., & Peterson, C. G. (1990). Effects of light and wave disturbance on vertical zonation of attached microalgae in a large reservoir. *Journal of Phycology*, 26(3), 450-457.  
doi:10.1111/j.0022-3646.1990.00450.x

Holland, R. S., & Hergenrader, G. L. (1981). Bacterial epibionts of diaptomid copepods. *Transactions of the American Microscopical Society*, 100(1), 56-65.

Holland, R. S., & Hergenrader, G. L. (1981). *Rhopalosolen saccatus* Fott: An epizoophyte from an alkaline lake in Nebraska. [Rhopalosolen saccatus (Fott): An Epizoophyte of Zooplankton from a Sandhills Lake] *American Midland Naturalist*, 106(2), 403-405.

Hornbeck, G. (2000). Ogallala trout. *Outdoor Life*, 205(5), 112-113.

Jameson, M. L. (1989). Diversity of coprophagous scarabaeidae (coleoptera) in grazed versus ungrazed sandhills prairie in western Nebraska. *Transactions of the Nebraska Academy of Sciences*, 17, 29-35.

Janovy Jr, J., Clopton, R. E., Clopton, D. A., Snyder, S. D., Efting, A., & Krebs, L. (1995). Species density distributions as null models for ecologically significant interactions of parasite species in an assemblage. *Ecological Modelling*, 77(2), 189-196.

Janovy Jr, J., & Kutish, G. W. (1988). A model of encounters between host and parasite populations. *Journal of Theoretical Biology*, 134(3), 391-401.

Janovy Jr, J., McDowell, M. A., & Ferdig, M. T. (1991). The niche of *Salsuginus thalkeni*, a gill parasite of *Fundulus zebrinus*. *The Journal of Parasitology*, 77(5), 697-702.

Janovy Jr, J., Snyder, S. D., & Clopton, R. E. (1997). Evolutionary constraints on population structure: The parasites of *Fundulus zebrinus* (pisces: Cyprinodontidae) in the South Platte River of Nebraska. *The Journal of Parasitology*, 83(4), 584-592.

Janovy Jr, J. (2002). Concurrent infections and the community ecology of helminth parasites. [Defining the field: Concurrent infections and the community ecology of helminth parasites] *Journal of Parasitology*, 88(3), 440-445. doi:10.1645/0022-3395(2002)088[0440:CIATCE]2.0.CO;2

Janovy Jr, J., Clopton, R. E., & Percival, T. J. (1992). The roles of ecological and evolutionary influences in providing structure to parasite species assemblages: Parasite biogeography and coevolution. *The Journal of Parasitology*, 78(4), 630-640.

Janovy Jr, J., & Hardin, E. L. (1987). Population dynamics of the parasites in *Fundulus zebrinus* in the Platte River of Nebraska. *The Journal of Parasitology*, 73(4), 689-696.

JANOVY JR, J., & HARDIN, E. L. (1988). Diversity of the parasite assemblage of *Fundulus zebrinus* in the Platte River of Nebraska. *The Journal of Parasitology*, 74(2), 207-213.

JANOVY JR, J., RUHNKE, T. R., & WHEELER, T. A. (1989). *Salsuginus thalkeni* n. sp.(monogenea: Ancyrocephalidae) from *Fundulus zebrinus* in the South Platte River of Nebraska. *The Journal of Parasitology*, 75(3), 344-347.

Janovy Jr., J., & Major, K. M. (2009). Why we have field stations: Reflections on the cultivation of biologists. *Bioscience*, 59(3), 217-222.

Janovy, J., Ferdig, M. T., & McDowell, M. A. (1990). A model of dynamic behavior of a parasite species assemblage. *Journal of Theoretical Biology*, 142(4), 517-529.

Joern, A. (1986). Temporal constancy in grasshopper assemblies (orthoptera: Acrididae). *Ecological Entomology*, 11(4), 379-385.

Joern, A., & Lawlor, L. R. (1981). Guild structure in grasshopper assemblages based on food and microhabitat resources. *Oikos*, 37, 93-104.

JOERN, A. (1981). Importance of behavior and coloration in the control of body temperature by *Brachystola magna* Girard (orthoptera: Acrididae). *Acrida*, 10(3), 117-130.

Joern, A. (1982). Distributions, densities, and relative abundances of grasshoppers (orthoptera: Acrididae) in a Nebraska sandhills prairie. *Prairie Naturalist*, 14(2), 37-45.

Joern, A. (1982). Vegetation structure and microhabitat selection in grasshoppers (orthoptera, Acrididae). *The Southwestern Naturalist*, 27(2), 197-209.

Joern, A. (1983). Host plant utilization by grasshoppers (orthoptera: Acrididae) from a sandhills prairie. *Journal of Range Management*, 36(6), 793-797.

Joern, A. (1983). Small-scale displacements of grasshoppers (orthoptera: Acrididae) within arid grasslands. *Journal of the Kansas Entomological Society*, 56(2), 131-139.

Joern, A. (1985). Grasshopper dietary (orthoptera: Acrididae) from a Nebraska sand hills prairie. *Transactions of the Nebraska Academy of Sciences*, 13, 21-32.

Joern, A. (1986). Experimental study of avian predation on coexisting grasshopper populations (orthoptera: Acrididae) in a sandhills grassland. *Oikos*, 46, 243-249.

JOERN, A. (1988). Foraging behavior and switching by the grasshopper sparrow *Ammodramus savannarum* searching for multiple prey in a heterogeneous environment. *The American Midland Naturalist*, 119(2), 225-234.

Joern, A. (1992). Variable impact of avian predation on grasshopper assemblies in sandhills grassland. *Oikos*, 64(3), 458-463.

Joern, A. (2002). Context-dependent foraging and enemy-free space: Grasshopper sparrows (*Ammodramus savannarum*) searching for grasshoppers (Acrididae). *Ecoscience*, 9(2), 231-240.

JOERN, A., & ALWARD, R. (1988). Effect of nitrogen fertilization on choice among grasses by the grasshopper *Phoetaliotes nebrascensis* (orthoptera: Acrididae). *Annals of the Entomological Society of America*, 81(2), 240-244.

Joern, A., & Behmer, S. T. (1997). Importance of dietary nitrogen and carbohydrates to survival, growth, and reproduction in adults of the grasshopper *Ageneotettix deorum* (orthoptera: Acrididae). *Oecologia*, 112(2), 201-208.

Joern, A., & Behmer, S. T. (1998). Impact of diet quality on demographic attributes in adult grasshoppers and the nitrogen limitation hypothesis. *Ecological Entomology*, 23(2), 174-184. doi:10.1046/j.1365-2311.1998.00112.x

Joern, A., Danner, B. J., Logan, J. D., & Wolessensky, W. (2006). Natural history of mass-action in predator-prey models: A case study from wolf spiders and grasshoppers. *The American Midland Naturalist*, 156(1), 52-64. doi:10.1674/0003-0031(2006)156[52:NHOMIP]2.0.CO;2

Joern, A., & Hoagland, K. D. (1996). In defense of whole-community bioassays for risk assessment. *Environmental Toxicology and Chemistry*, 15(4), 407-409. doi:10.1002/etc.5620150401

Joern, A., & Klucas, G. (1993). Intra-and interspecific competition in adults of two abundant grasshoppers (orthoptera: Acrididae) from a sandhills grassland. *Environmental Entomology*, 22(2), 352-361.

Joern, A., Mitschler, R., & O'Leary, H. (1986). Activity and time budgets of three grasshopper species (orthoptera: Acrididae) from a sandhills grassland. [Activity and time budgets in common sandhills prairie grasshoppers] *Journal of the Kansas Entomological Society*, 59(1), 1-6.

Joern, A., & Mole, S. (2005). The plant stress hypothesis and variable responses by blue grama grass (*Bouteloua gracilis*) to water, mineral nitrogen, and insect herbivory. *Journal of Chemical Ecology*, 31(9), 2069-2090. doi:10.1007/s10886-005-6078-3

Joern, A., Provin, T., & Behmer, S. T. (2012). Not just the usual suspects: Insect herbivore populations and communities are associated with multiple plant nutrients. *Ecology*, 93(5), 1002-1015. doi:10.1890/11-1142.1

Joern, A., & Rudd, N. T. (1982). Impact of predation by the robber fly *Proctacanthus milbertii* (diptera: Asilidae) on grasshopper (orthoptera: Acrididae) populations. *Oecologia*, 55(1), 42-46.

Johnsgard, P. A. (1990). Additional observations on the birds of the lake McConaughy region. *Nebraska Bird Review*, 58(2), 52-4.

Johnsgard, P. A. (1998). Endemicity and regional biodiversity in Nebraska's breeding avifauna. *Nebraska Bird Review*, 66(4), 115-121.

Jones, S. M., Ballinger, R. E., & Porter, W. P. (1987). Physiological and environmental sources of variation in reproduction: Prairie lizards in a food rich environment. *Oikos*, 48(3), 325-335.

Jones, S. M., Waldschmidt, S. R., & Potvin, M. A. (1987). An experimental manipulation of food and water: Growth and time-space utilization of hatchling lizards (*Sceloporus undulatus*). *Oecologia*, 73(1), 53-59.

Jones, S. M., & Ballinger, R. E. (1987). Comparative life histories of *Holbrookia maculata* and *Sceloporus undulatus* in western Nebraska. *Ecology*, 68(6), 1828-1838.

Jones, S. M., & Droege, D. L. (1980). Home range size and spatial distributions of two sympatric lizard species (*Sceloporus undulatus*, *Holbrookia maculata*) in the sand hills of Nebraska. *Herpetologica*, 36(2), 127-132.

Jordon-Thaden, I., & Louda, S. M. (2003). Chemistry of cirsium and carduus: A role in ecological risk assessment for biological control of weeds? [Chemistry of thistles: A role in ecological risk assessment for biological control of weeds?] *Biochemical Systematics and Ecology*, 31(12), 1353-1396.

Jorgensen, J. G., & Paseka, J. (2001). 1999 (eleventh) report of the NOU records committee. *Nebraska Bird Review*, 69(2).

Jorgensen, J. G., & Paseka, J. (2002). 2002 (12th) report of the NOU records committee. *Nebraska Bird Review*, 70(2).

Jurgensen, T. A., & Hoagland, K. D. (1990). Effects of short-term pulses of atrazine on attached algal communities in a small stream. *Archives of Environmental Contamination and Toxicology*, 19(4), 617-623.

Kapustka, L. A., Arnold, P. T., & Lattimore, P. T. (1985). Interactive responses of associative diazotrophs from a Nebraska sand hills grassland. *Special Publication... of the British Ecological Society*, 4, 149-158.

Kapustka, L. A., & DuBois, J. D. (1982). Legume distribution and nodulation in Arapaho Prairie, Arthur County, Nebraska. *Prairie Naturalist*, 14(1), 1-5.

Kapustka, L. A., & Dubois, J. D. (1987). Dinitrogen fixation by cyanobacteria and associative rhizosphere bacteria in the Arapho Prairie in the sand hills of Nebraska. *American Journal of Botany*, 74(1), 107-113.

Kapustka, L. A., Morrison, L. C., & DuBois, J. D. (1988). Dinitrogen fixation in the wet meadows and emergent zones of two Nebraska sandhills lakes. *American Midland Naturalist*, 120(2), 398-404.

Kaspari, M. (1990). Prey preparation and the determinants of handling time. *Animal Behaviour*, 40(1), 118-126.

Kaspari, M. (1991). Central place foraging in grasshopper sparrows: Opportunism or optimal foraging in a variable environment? *Oikos*, 60(3), 307-312.

Kaspari, M. (1991). Prey preparation as a way that grasshopper sparrows (*Ammodramus savannarum*) increase the nutrient concentration of their prey. [Preparation as a strategy to maximize nutrient concentration in prey] *Behavioral Ecology*, 2(3), 234-241.

doi:10.1093/beheco/2.3.234

Kaspari, M., & Joern, A. (1993). Prey choice by three insectivorous grassland birds: Reevaluating opportunism. *Oikos*, 68(3), 414-430.

Kaspari, M., & O'Leary, H. (1988). Nonparental attendants in a north-temperate migrant. *Auk*, 105(4), 792-793.

Kaufmann, S. A., & Lynch, J. D. (1991). Courtship, eggs, and development of the plains topminnow in Nebraska (actinopterygii: Fundulidae). *Prairie Naturalist*, 23(1), 41-45.

KAUL, R. B. (1993). Meristic and organogenetic variation in *Ruppia occidentalis* and *R. maritima*. *International Journal of Plant Sciences*, 154(3), 416-424.

Kaul, R. B., Rolfsmeier, S. B., & Esch, J. J. (1991). The distribution and reproductive phenology of the milkweeds (asclepiadaceae: *Asclepias* and *cynanchum*) in Nebraska. *Transactions of the Nebraska Academy of Sciences*, 18, 127-140.

Kaul, R. B., & Rolfsmeier, S. B. (1994). Diversity of vascular plants and intensity of plant collecting in Nebraska. *Transactions of the Nebraska Academy of Sciences*, 21, 13-20.

KEELER, K. H. (1987). Survivorship and fecundity of the polycarpic perennial *mentzelia nuda* (loasaceae) in Nebraska sandhills prairie. *American Journal of Botany*, 74(6), 785-791.

Keeler, K. H. (1993). Fifteen years of colony dynamics in *Pogonomyrmex occidentalis*, the western harvester ant, in western Nebraska. *The Southwestern Naturalist*, 38(3), 286-289.

Keeler, K. H. (1979). Nocturnal pollination of *Abronia fragrans* (nyctaginaceae). *Southwestern Naturalist*, 24, 692-693.

Keeler, K. H. (1979). Species with extrafloral nectaries in a temperate flora (Nebraska). *Prairie Naturalist*, 11(1), 33-38.

Keeler, K. H. (1980). Distribution of plants with extrafloral nectaries in temperate communities. *American Midland Naturalist*, 104(2), 274-280.

Keeler, K. H. (1980). The extrafloral nectaries of *Ipomoea leptophylla* (convolvulaceae). *American Journal of Botany*, 67(2), 216-222.

Keeler, K. H. (1981). Function of *Mentzelia nuda* (loasaceae) postfloral nectaries in seed defense. *American Journal of Botany*, 68(2), 295-299.

Keeler, K. H. (1981). A model of selection for facultative nonsymbiotic mutualism. *American Naturalist*, 118(4), 488-498.

Keeler, K. H. (1982). Preliminary report of colony survivorship in the western harvester ant (*Pogonomyrmex occidentalis*) in western Nebraska. *Southwestern Naturalist*, 27(2), 245-246.

Keeler, K. H. (1988). Colony survivorship in *Pogonomyrmex occidentalis*, western harvester ant, in western Nebraska. *The Southwestern Naturalist*, 33(4), 480-482.

Keeler, K. H. (1991). Survivorship and recruitment in a long-lived prairie perennial, *Ipomoea leptophylla* (convolvulaceae). *American Midland Naturalist*, 126(1), 44-60.

Keeler, K. H., Harrison, A. T., & Vescio, L. S. (1980). The flora and sandhills prairie communities of Arapaho Prairie, Arthur County, Nebraska. *Prairie Naturalist*, 12, 65-78.

Keeler, K. H., & Kaul, R. B. (1979). Morphology and distribution of petiolar nectaries in *Ipomoea* (convolvulaceae). *American Journal of Botany*, 66(8), 946-952.

Keeler, K. H., & Kaul, R. B. (1984). Distribution of defense nectaries in *Ipomoea* (convolvulaceae). *American Journal of Botany*, 71(10), 1364-1372.

Kinney, T. A., & Lynch, J. D. (1991). The fecundity and reproductive season of *Fundulus sciadicus* in Nebraska (actinopterygii: Fundulidae). *Transactions of the Nebraska Academy of Sciences*, 18, 101-104.

Knight, S. A., Janovy Jr, J., & Current, W. L. (1977). *Myxosoma funduli* kudo 1918 (protozoa: Myxosporida) in *Fundulus kansasae*: Summer epizootiology. *The Journal of Parasitology*, 63(5), 897-902.

Knight, S. A., Janovy Jr, J., & Current, W. L. (1980). *Myxosoma funduli* kudo 1918 (protozoa: Myxosporida) in *Fundulus kansasae* (pisces: Cyprinodontidae): Annual prevalence and geographic distribution. *The Journal of Parasitology*, 66(5), 806-810.

Kren, J., & Zoerb, A. C. (1997). Northern wheatear (*Oenanthe oenanthe*). *The Birds of North America*, (316), 24.

Krupa, J. J. (2000). The importance of naturalists as teachers & the use of natural history as a teaching tool. *The American Biology Teacher*, 62(8), 553-558.

Kruse, K. C., & Francis, M. G. (1977). A predation deterrent in larvae of the bullfrog, *Rana catesbeiana*. *Transactions of the American Fisheries Society*, 106(3), 248-252.  
doi:10.1577/1548-8659(1977)106<248:APDILO>2.0.CO;2

Kugler, K. A., & Geluso, K. (2009). Distribution of the eastern woodrat (*Neotoma floridana campestris*) in southern Nebraska. *Western North American Naturalist*, 69(2), 175-179.  
doi:10.3398/064.069.0205

Kula, R. R., & Clopton, R. E. (1999). *Amoebogregarina nigra* n. gen., n. comb.(apicomplexa: Gregarinidae) from adult *Melanoplus differentialis* (orthoptera: Acrididae) in southeastern Nebraska. *The Journal of Parasitology*, 85(2), 321-325.

Lamp, W. O., & McCarty, M. K. (1982). Biology of predispersal seed predators of the platte thistle, *Cirsium canescens*. *Journal of the Kansas Entomological Society*, 55(2), 305-316.

Lamp, W. O., & McCarty, M. K. (1982). Predispersal seed predation of a native thistle, *Cirsium canescens*. *Environmental Entomology*, 11(4), 847-851.

Lamp, W. O., & McCarty, M. K. (1981). Biology and ecology of the platte thistle (*Cirsium canescens*). *Weed Science*, 29(6), 686-692.

Lamp, W. O., & McCarty, M. K. (1982). Observations of *Corythucha distincta* (hemiptera: Tingidae) on platte thistle in Nebraska. *Journal of the Kansas Entomological Society*, 55(1), 34-36.

Langford, G. J., & Janovy Jr, J. (2009). Comparative life cycles and life histories of North American *Rhabdias* spp.(nematoda: Rhabdiasidae): Lungworms from snakes and anurans. *Journal of Parasitology*, 95(5), 1145-1155. doi:10.1645/GE-2044.1

Langford, G. J., & Janovy Jr, J. (2013). Host specificity of North American *Rhabdias* spp. (nematoda: Rhabdiasidae): Combining field data and experimental infections with a molecular phylogeny. *The Journal of Parasitology*, 99(2), 277-286. doi:10.1645/GE-3217.1

Ledder, G., Logan, J. D., & Joern, A. (2004). Dynamic energy budget models with size-dependent hazard rates. *Journal of Mathematical Biology*, 48(6), 605-622.  
doi:10.1007/s00285-003-0263-1

Lemen, C. A., & Freeman, P. W. (1985). Tracking mammals with fluorescent pigments: A new technique. *Journal of Mammalogy*, 66(1), 134-136.

Lewis, R. J., Jensen, S. I., DeNicola, D. M., Miller, V. I., Hoagland, K. D., & Ernst, S. G. (1997). Genetic variation in the diatom *Fragilaria capucina* (*fragilariaeae*) along a latitudinal gradient across North America. *Plant Systematics and Evolution*, 204(1-2), 99-108.

Lind, E. M., Borer, E., Seabloom, E., Adler, P., Bakker, J. D., Blumenthal, D. M., . . . Wragg, P. D. (2013). Life-history constraints in grassland plant species: A growth-defence trade-off is the norm. *Ecology Letters*, 16(4), 513-521.

Lipar, J. L., Ketterson, E. D., Nolan, V., & Casto, J. M. (1999). Egg yolk layers vary in the concentration of steroid hormones in two avian species. *General and Comparative Endocrinology*, 115(2), 220-227.

Lipar, J. L., & Ketterson, E. D. (2000). Maternally derived yolk testosterone enhances the development of the hatching muscle in the red-winged blackbird *Agelaius phoeniceus*. *Proceedings of the Royal Society of London Series B: Biological Sciences*, 267(1456), 2005-2010.

Logan, J. D., Joern, A., & Wolesensky, W. (2003). Chemical reactor models of optimal digestion efficiency with constant foraging costs. *Ecological Modelling*, 168(1), 25-38.

Logan, J. D., Joern, A., & Wolesensky, W. (2004). Control of CNP homeostasis in herbivore consumers through differential assimilation. *Bulletin of Mathematical Biology*, 66(4), 707-725. doi:10.1016/j.bulm.2003.10.008

Logan, J. D., Joern, A., & Wolesensky, W. (2004). Mathematical model of consumer homeostasis control in plant-herbivore dynamics. *Mathematical and Computer Modelling*, 40(3), 447-456.

Logan, J. D., Wolesensky, W., & Joern, A. (2006). Temperature-dependent phenology and predation in arthropod systems. *Ecological Modelling*, 196(3), 471-482.

Logan, J. D., Janovy, J., & Bunker, B. E. (2012). The life cycle and fitness domain of gregarine (apicomplexa) parasites. *Ecological Modelling*, 233, 31-40.

Logan, J. D., Wolesensky, W., & Joern, A. (2007). Insect development under predation risk variable temperature, and variable food quality. *Mathematical Biosciences and Engineering: MBE*, 4(1), 47-65.

Louda, S. M. (1989). Differential predation pressure: A general mechanism for structuring plant communities along complex environmental gradients? *Trends in Ecology & Evolution*, 4(6), 158-159.

Louda, S. M. (1997). Ecological effects of a host range expansion by a weevil imported for the biological control of weeds. *Bulletin of the Ecological Society of America*, 78, 122.

Louda, S. M. (1997). Impacts of native and non-native insect natural enemies on the dynamics of herbaceous perennial plants. *Bulletin of the Ecological Society of America*, 78, 43.

Louda, S. M. (1999). Ecology of interactions needed in biological control practice and policy.

*Bulletin of the British Ecological Society*, 29(4), 834-841.

Louda, S. M., Arnett, A. E., Rand, T. A., & Russell, F. L. (2003). Invasiveness of some biological control insects and adequacy of their ecological risk assessment and regulation: Population biology of invasive species. *Conservation Biology*, 17(1), 73-82.  
doi:10.1046/j.1523-1739.2003.02020.x

Louda, S. M., Potvin, M. A., & Collinge, S. K. (1990). Predispersal seed predation, postdispersal seed predation and competition in the recruitment of seedlings of a native thistle in sandhills prairie. *American Midland Naturalist*, 124(1), 105-113.

Louda, S. M., & Rodman, J. E. (1996). Insect herbivory as a major factor in the shade distribution of a native crucifer (*Cardamine cordifolia* A. gray, bittercress). *Journal of Ecology*, 84(2), 229-237.

Louda, S. M., Russell, F. L., & Rand, T. A. (2003). Caution needed in biological control efforts. *Conservation in Practice*, 4, 45-46.

Louda, S. M., Simberloff, D., Boettner, G., Connor, J., Kendall, D., & Arnett, A. E. (1998). Insights from data on the nontarget effects of the flowerhead weevil. *Biocontrol News and Information*, 19, 70N-72N.

Louda, S. M. (1998). Population growth of *Rhinocyllus conicus* (coleoptera: Curculionidae) on two species of native thistles in prairie. *Environmental Entomology*, 27(4), 834-841.

Louda, S. M., Kendall, D., Connor, J., & Simberloff, D. (1997). Ecological effects of an insect introduced for the biological control of weeds. *Science*, 277(5329), 1088-1090.  
doi:10.1126/science.277.5329.1088

Louda, S. M., & Masters, R. A. (1993). Biological control of weeds in great plains rangelands. *Great Plains Research: A Journal of Natural and Social Sciences*, 3(1), 125-247.

LOUDA, S. M., & O'BRIEN, C.,W. (2002). Unexpected ecological effects of distributing the exotic weevil, *Larinus planus* (F.), for the biological control of Canada thistle. *Conservation Biology*, 16(3), 717-727. doi:10.1046/j.1523-1739.2002.00541.x

Louda, S. M., Pemberton, R. W., Johnson, M. T., & Follett, P. A. (2003). Nontarget effects-the achilles' heel of biological control? Retrospective analyses to reduce risk associated with biocontrol introductions. *Annual Review of Entomology*, 48(1), 365-396.

Louda, S. M., & Potvin, M. A. (1995). Effect of inflorescence-feeding insects on the demography and lifetime of a native plant. *Ecology*, 76(1), 229-245. doi:10.2307/1940645

Louda, S. M., Rand, T. A., Arnett, A. E., McClay, A. S., Shea, K., & McEachern, A. K. (2005). Evaluation of ecological risk to populations of a threatened plant from an invasive biocontrol insect. *Ecological Applications*, 15(1), 234-249. doi:10.1890/03-5212

Louda, S. M., Rand, T. A., Kula, A., Arnett, A. E., West, N. M., & Tenhumberg, B. (2011). Priority resource access mediates competitive intensity between an invasive weevil and native floral herbivores. *Biological Invasions*, 13(10), 2233-2248. doi:10.1007/s10530-011-0036-5

Louda, S. M., Rand, T. A., Russell, F. L., & Arnett, A. E. (2005). Assessment of ecological risks in weed biocontrol: Input from retrospective ecological analyses. *Biological Control*, 35(3), 253-264.

Louda, S. M., & Stiling, P. (2004). The double- • Edged sword of biological control in conservation and restoration. *Conservation Biology*, 18(1), 50-53. doi:10.1111/j.1523-1739.2004.00070.x

Lynch, J. D. (1988). Introduction, establishment, and dispersal of western mosquitofish in Nebraska (actinopterygii: Poeciliidae). *Prairie Nat*, 20, 203-216.

Lynch, J. D. (1978). The distribution of leopard frogs (*Rana blairi* and *Rana pipiens*)(amphibia, anura, ranidae) in Nebraska. *Journal of Herpetology*, 12(2), 157-162.

Lynch, J. D. (1985). Annotated checklist of the amphibians and reptiles of Nebraska. *Transactions of the Nebraska Academy of Sciences*, 13, 33-57.

Lynch, J. D. (1988). Habitat utilization by an introduced fish, *Gambusia affinis*, in Nebraska (actinopterygii: Poeciliidae). [Habitat utilization by an introduced minnow, *Gambusia affinis*, in Nebraska (Actinopterygii: Poeciliidae)] *Transactions of the Nebraska Academy of Sciences*, 16, 63-67.

Lynch, J. D., & Roh, B. R. (1996). An ichthyological survey of the forks of the Platte River in western Nebraska. *Transactions of the Nebraska Academy of Sciences*, 23, 65-84.

Marion, K. R., Bizer, J. R., & Sexton, O. J. (1979). A between-clutch comparison of hatchling weights in the lizard *Sceloporus undulatus* (reptilia: Squamata: Lacertilia). *Herpetologica*, 35(2), 111-114.

Maron, J. L., Combs, J. K., & Louda, S. M. (2002). Insect herbivory on thistles: Congruent interaction strengths in coastal vs. continental dunes. *Ecology*, 83, 3382-3392.

Matthews, R. W., & Matthews, J. R. (2005). Biological notes on ground-nesting digger wasps from western Nebraska. *Prairie Naturalist*, 37(2), 61-71.

MCDOWELL, M. A., FERDIG, M. T., & JANOVY JR, J. (1992). Dynamics of the parasite assemblage of *Pimephales promelas* in Nebraska. *The Journal of Parasitology*, 78(5), 830-836.

McKenzie, D. A. (2005). Plant composition of mixed-grass prairie of Cedar Point Biological Research Station, Keith County, Nebraska. *Transactions of the Nebraska Academy of Sciences*, 30, 17-23.

McLennan, D. A. (1993). Temporal changes in the structure of the male nuptial signal in the brook stickleback, *Culaea inconstans* (kirtland). *Canadian Journal of Zoology*, 71(6), 1111-1119. doi:10.1139/z93-151

McLennan, D. A. (1994). Changes in female colour across the ovulatory cycle in the brook stickleback, *Culaea inconstans* (kirtland). *Canadian Journal of Zoology*, 72(1), 144-153. doi:10.1139/z94-018

Messenger, C. (1997). The sphinx moths (lepidoptera: Sphingidae) of Nebraska. *Transactions of the Nebraska Academy of Sciences*, 24, 89-141.

Mole, S., & Joern, A. (1993). Foliar phenolics of Nebraska sandhills prairie graminoids: Between-years, seasonal, and interspecific variation. *Journal of Chemical Ecology*, 19(9), 1861-1874.

Mole, S., & Joern, A. (1994). Feeding behavior of graminivorous grasshoppers in response to host-plant extracts, alkaloids, and tannins. *Journal of Chemical Ecology*, 20(12), 3097-3109.

Mole, S., Joern, A., O'Leary, M., H., & Madhavan, S. (1994). Spatial and temporal variation in carbon isotope discrimination in prairie graminoids. *Oecologia*, 97(3), 316-321.

Møller, A. P., Merino, S., Brown, C. R., & Robertson, R. J. (2001). Immune defense and host sociality: A comparative study of swallows and martins. *The American Naturalist*, 158(2), 136-145. doi:10.1086/321308

Montanucci, R. R. (1979). Notes on systematics of horned lizards allied to *Phrynosoma orbiculare* (lacertilia: Iguanidae). *Herpetologica*, 35(2), 116-124.

Moody-Weis, J., Antonovics, J., Alexander, H. M., & Pilson, D. (2008). Predicting local colonization and extinction dynamics from coarser-scale surveys. *Ecography*, 31(1), 61-72. doi:10.1111/j.2008.0906-7590.05282.x

Moore, A. T., Edwards, E. A., Bomberger Brown, M., Komar, N., & Brown, C. R. (2007). Ecological correlates of buggy creek virus infection in *Oeciacus vicarius*, southwestern

Nebraska, 2004. *Journal of Medical Entomology*, 44(1), 42-49. doi:10.1603/0022-2585(2007)44[42:ECOBCV]2.0.CO;2

Morrison, L. C., DuBois, J. D., & Kapustka, L. A. (1986). The vegetational response of a Nebraska sandhills grassland to a naturally occurring fall burn. *Prairie Naturalist*, 18(3), 179-184.

Morrison, R., & Powell, L. (1988). New distributional records of lizards in Wyoming. *Transactions of the Nebraska Academy of Sciences*, 16, 85-86.

Moser, W. A. (1991). Leeches (annelida: Hirudinea) in central and western Nebraska. *Transactions of the Nebraska Academy of Sciences*, 18, 87-92.

Nietfeldt, J. W., Jones, S. M., Droke, D. L., & Ballinger, R. E. (1980). Rate of thermal acclimation in larval *Ambystoma tigrinum*. *Journal of Herpetology*, 14(3), 209-211.

Niewiarowski, P. H., & Roosenburg, W. (1993). Reciprocal transplant reveals sources of variation in growth rates of the lizard *Sceloporus undulatus*. *Ecology*, 74, 1992-2002.  
doi:10.2307/1940842

O'Brien, V., & Brown, C. R. (2010). Group size and nest spacing affect buggy creek virus (togaviridae: Alphavirus) infection in nestling house sparrows. *PLoS One*, 6(9), e25521-e25521. doi:10.1371/journal.pone.0025521

O'Brien, V., Meteyer, C. U., Ip, H. S., Long, R. R., & Brown, C. R. (2010). Pathology and virus detection in tissues of nestling house sparrows naturally infected with buggy creek virus (togaviridae). *Journal of Wildlife Diseases*, 46(1), 23-32.

O'Brien, V.,A., Meteyer, C. U., Reisen, W. K., Ip, H. S., & Brown, C. R. (2010). Prevalence and pathology of west nile virus in naturally infected house sparrows, western Nebraska, 2008.

*The American Journal of Tropical Medicine and Hygiene*, 82(5), 937-944.

doi:10.4269/ajtmh.2010.09-0515

O'Brien, V.,A., Moore, A. T., Huyvaert, K. P., & Brown, C. R. (2008). No evidence for spring re-introduction of an arbovirus by cliff swallows. *The Wilson Journal of Ornithology*, 120(4), 910-913. doi:10.1676/08-028.1

O'Brien, V.,A., Moore, A. T., Young, G. R., Komar, N., Reisen, W. K., & Brown, C. R. (2011).

An enzootic vector-borne virus is amplified at epizootic levels by an invasive avian host.

*Proceedings of the Royal Society B: Biological Sciences*, 278(1703), 239-246.

doi:10.1098/rspb.2010.1098

O'Brien Valerie, A., & Brown Charles, R. (2012). Arbovirus infection is a major determinant of fitness in house sparrows (*Passer domesticus*) that invade cliff swallow (*Petrochelidon pyrrhonota*) colonies. *The Auk*, 129(4), 707-715. doi:10.1525/auk.2012.12036

Oedekoven, M. A., & Joern, A. (1998). Stage-based mortality of grassland grasshoppers (acrididae) from wandering spider (lycosidae) predation. *Acta Oecologica*, 19(6), 507-515.

Oedekoven, M. A., & Joern, A. (2000). Plant quality and spider predation affects grasshoppers (acrididae): Food-quality-dependent compensatory mortality. [Host plant quality and risk to spider predation by grasshoppers (Acrididae): state-dependent compensatory mortality] *Ecology*, 81(1), 66-77. doi:10.1890/0012-9658(2000)081[0066:PQASPA]2.0.CO;2

Okuyama, T. (2007). Prey of two species of jumping spiders in the field. *Applied Entomology and Zoology*, 42(4), 663-668. doi:10.1303/aez.2007.663

OLSON, P. D., & NICKOL, B. B. (1996). Comparison of *Leptorhynchoides thecatus* (acanthocephala) recruitment into green sunfish and largemouth bass populations. *The Journal of Parasitology*, 82(5), 702-706.

Padhi, A., Moore, A. T., Brown, M. B., Foster, J. E., Pfeffer, M., & Brown, C. R. (2011). Isolation by distance explains genetic structure of buggy creek virus, a bird-associated arbovirus. *Evolutionary Ecology*, 25(2), 403-416. doi:10.1007/s10682-010-9419-9

Padhi, A., Moore, A. T., Brown, M. B., Foster, J. E., Pfeffer, M., Gaines, K. P., . . . Brown, C. R. (2008). Phylogeographical structure and evolutionary history of two buggy creek virus lineages in the western great plains of North America. *Journal of General Virology*, 89(9), 2122-2131. doi:10.1099/vir.0.2008/001719-0

Pappas, P. W. (1978). Tryptic and protease activities in the normal and *Hymenolepis diminuta*-infected rat small intestine. *The Journal of Parasitology*, 64(3), 562-564.

Parrott, A., & David Logan, J. (2010). Effects of temperature variation on TSD in turtle (*C. picta*) populations. *Ecological Modelling*, 221(10), 1378-1393.

Paseka, J. (1997). Survey of bald and golden eagles in Nebraska. *Nebraska Bird Review*, 54(1), 36-40.

Paseka, J., Peyton, M. M., & Knaggs, R. (1997). Bald eagle counts at two central Nebraska public power and irrigation district facilities. *Nebraska Bird Review*, 65(1), 40-41.

Paulsen, M. J. (2002). Observations on possible myrmecophily in *Stephanucha pilipennis* kraatz (coleoptera: Scarabaeidae: Cetoniinae) in western Nebraska. *The Coleopterists Bulletin*, 56(3), 451-452. doi:10.1649/0010-065X(2002)056[0451:OOPMIS]2.0.CO;2

Paulsen, M. J. (2006). A new species and new records of *aphodius* illiger (coleoptera: Scarabaeidae: Aphodiinae) from mammal burrows in Nebraska. *Insecta Mundi*, 20(1-2), 95-100.

Percival Cook, T.,J., Janovy Jr, J., & Clopton, R. E. (2001). Epimerite-host epithelium relationships among eugregarines parasitizing the damselflies *Enallagma civile* and *Ischnura verticalis*. *Journal of Parasitology*, 87(5), 988-996. doi:10.1645/0022-3395(2001)087[0988:EHERAE]2.0.CO;2

Percival, T. J., Clopton, R. E., & Janovy, J. (1995). Two new menosporine gregarines, *Hoplorhynchus acanthatholius* N. sp. and *Steganorhynchus dunwoodyi* N. G., N. sp. (apicomplexa: Eugregarinorida: Actinocephalidae) from coenagrionid damselflies (odonata: Zygoptera). *Journal of Eukaryotic Microbiology*, 42(4), 406-410. doi:10.1111/j.1550-7408.1995.tb01603.x

Peterson, C. G., & Hoagland, K. D. (1990). Effects of wind-induced turbulence and algal mat development on epilithic diatom succession in a large reservoir. *Archiv Für Hydrobiologie*, 118(1), 47-68.

Peterson, C. G., Hoagland, K. D., & Stevenson, R. J. (1990). Timing of wave disturbance and the resistance and recovery of a freshwater epilithic microalgal community. *Journal of the North American Benthological Society*, 9(1), 54-67.

Peterson, C. G., Hoagland, K. D., & Stevenson, R. J. (1990). Timing of wave disturbance and the resistance and recovery of a freshwater epilithic microalgal community. [Resistance and recovery in freshwater epilithic microalgal communities: importance of timing of wave disturbance] *Journal of the North American Benthological Society*, 9(1), 54-67.

Peyton, M. M., & Paseka, J. (2004). Lake McConaughy and Nebraska piping plover recovery goals. *Nebraska Bird Review*, 72(4), 148-152.

Pfeffer, M., Foster, J. E., Edwards, E. A., Brown, M. B., Komar, N., & Brown, C. R. (2006). Phylogenetic analysis of buggy creek virus: Evidence for multiple clades in the western great plains, united states of america. *Applied and Environmental Microbiology*, 72(11), 6886-6893. doi:10.1128/AEM.00868-06

Pilson, D. (1999). Plant hybrid zones and insect host range expansion. *Ecology*, 80(2), 407-415. doi:10.1890/0012-9658(1999)080[0407:PHZAIH]2.0.CO;2

Pilson, D. (2000). Herbivory and natural selection on flowering phenology in wild sunflower, *Helianthus annuus*. [Flowering phenology and resistance to herbivory in wild sunflower, *Helianthus annuus*] *Oecologia*, 122(1), 72-82. doi:10.1007/PL00008838

PILSON, D., & DECKER, K. L. (2002). Compensation for herbivory in wild sunflower: Response to simulated damage by the head-clipping weevil. *Ecology*, 83(11), 3097-3107. doi:10.1890/0012-9658(2002)083[3097:CFHIWS]2.0.CO;2

Porath, M. T., & Peters, E. J. (1997). Walleye prey selection in Lake McConaughy, Nebraska: A comparison between stomach content analysis and feeding experiments. *Journal of Freshwater Ecology*, 12(4), 511-520. doi:10.1080/02705060.1997.9663565

Potthoff, T. L., & Lynch, J. D. (1986). Interpopulation variability in mouthparts in *Scaphiopus bombifrons* in Nebraska (amphibia: Pelobatidae). *Prairie Naturalist*, 18, 15-22.

Potvin, M. A. (1988). Seed rain on a Nebraska sandhills prairie. *Prairie Naturalist*, 20, 81-89.

Potvin, M. A. (1993). Establishment of native grass seedlings along a topographic/moisture gradient in the Nebraska sandhills. *The American Midland Naturalist*, 130(2), 248-261.

Potvin, M. A., & Harrison, A. T. (1984). Vegetation and litter changes of a Nebraska sandhills prairie protected from grazing. *Journal of Range Management*, 37(1), 55-58.

Price, T. D., Brown, C. R., & Brown, M. B. (2000). Evaluation of selection on cliff swallows. *Evolution*, 54(5), 1824-1827. doi:10.1111/j.0014-3820.2000.tb00727.x

Rand, T. A., & Louda, S. M. (2006). Spillover of agriculturally subsidized predators as a potential threat to native insect herbivores in fragmented landscapes. [Predator spillover across the agricultural-natural interface: implications for insects in fragmented landscapes] *Conservation Biology*, 20, 1720-1729. doi:10.1111/j.1523-1739.2006.00507.x

Rand, T. A., Russell, F. L., & Louda, S. M. (2004). Local-vs. landscape-scale indirect effects of an invasive weed on native plants 1. *Weed Technology*, 18, 1250-1254. doi:10.1614/0890-037X(2004)018[1250:LVLIEO]2.0.CO;2

Rand, T. A., & Louda, S. M. (2004). Exotic weed invasion increases the susceptibility of native plants to attack by a biocontrol herbivore. *Ecology*, 85(6), 1548-1554.

Rand, T. A., & Louda, S. M. (2006). Invasive insect abundance varies across the biogeographic distribution of a native host plant. *Ecological Applications*, 16(3), 877-890.  
doi:10.1890/1051-0761(2006)016[0877:IIAVAT]2.0.CO;2

Rannala, B. (1996). The sampling theory of neutral alleles in an island population of fluctuating size. *Theoretical Population Biology*, 50(1), 91-104.

Rannala, B. H., & Brown, C. R. (1994). Relatedness and conflict over optimal group size. *Trends in Ecology & Evolution*, 9(4), 117-119.

Rannala, B., & Hartigan, J. A. (1995). Identity by descent in island-mainland populations. *Genetics*, 139(1), 429-437.

Rannala, B., & Hartigan, J. A. (1996). Estimating gene flow in island populations. *Genetical Research*, 67(02), 147-158. doi:10.1017/S0016672300033607

Raouf, S. A., Smith, L. C., Brown, M. B., Wingfield, J. C., & Brown, C. R. (2006). Glucocorticoid hormone levels increase with group size and parasite load in cliff swallows. *Animal Behaviour*, 71(1), 39-48.

Rhoden, H. R., & Bolek, M. G. (2011). Distribution and reproductive strategies of *Gyrinicola batrachiensis* (oxyuroidea: Pharyngodonidae) in larvae of eight species of amphibians from Nebraska. *Journal of Parasitology*, 97(4), 629-635. doi:10.1645/GE-2670.1

Rhoden, H. R., & Bolek, M. G. (2012). Helminth and leech community structure in tadpoles and caudatan larvae of two amphibian species from western Nebraska. *Journal of Parasitology*, 98(2), 236-244. doi:10.1645/GE-2771.1

Rhodes, H. A., & Kondratieff, B. C. (1996). Annotated list of the stoneflies (plecoptera) of western Nebraska, USA. *Journal of the Kansas Entomological Society*, 69(2), 191-198.

Richardson, D. J. (2013). Helminth parasites of the raccoon (*Procyon lotor*), virginia opossum (*Didelphis virginiana*), and striped skunk (*Mephitis mephitis*) from Keith County, Nebraska. *Transactions of the Nebraska Academy of Sciences*, 33, 35-38.

Richardson, D. J., & Barger, M. A. (2005). Microhabitat specificity of *Macracanthorhynchus ingens* (acanthocephala: Oligacanthorhynchidae) in the raccoon (*Procyon lotor*). *Comparative Parasitology*, 72(2), 173-178. doi:10.1654/4199

Richardson, D. J., & Barger, M. A. (2006). Redescription of *Oligacanthorhynchus major* (Machado-Filho, 1963) Schmidt, 1972 (acanthocephala: Oligacanthorhynchidae) from the white-lipped peccary (*Tayassu pecari*) in Bolivia. *Comparative Parasitology*, 73(2), 157-160. doi:10.1654/4235.1

Richardson, S., & Janovy, J. (1990). *Actinocephalus carrilynnae* n. sp.(apicomplexa: Eugregarinorida) from the blue damselfly, *Enallagma civile* (Hagen). *Journal of Eukaryotic Microbiology*, 37(6), 567-570. doi:10.1111/j.1550-7408.1990.tb01266.x

Roche, E. A., Brown, C. R., & Brown, M. B. (2011). Heritable choice of colony size in cliff swallows: Does experience trump genetics in older birds? *Animal Behaviour*, 82(6), 1275-1285.

Roche, E. A., Brown, C. R., Brown, M. B., & Lear, K. M. (2013). Recapture heterogeneity in cliff swallows: Increased exposure to mist nets leads to net avoidance. *PloS One*, 8(3), e58092. doi:10.1371/journal.pone.0058092

Roemer, S. C., & Hoagland, K. D. (1979). Seasonal attenuation of quantum irradiance (400-700 NM) in three Nebraska reservoirs. *Hydrobiologia*, 63(1), 81-92.

Roemer, S. C., Hoagland, K. D., & Rosowski, J. R. (1984). Development of a freshwater periphyton community as influenced by diatom mucilages. *Canadian Journal of Botany*, 62(9), 1799-1813.

Rolfsmeier, S. B., Kaul, R. B., & Sutherland, D. M. (1987). New and corrected records of the flora of Nebraska. *Transactions of the Nebraska Academy of Sciences and Affiliated Societies*, 15, 49-52.

Rolfsmeier, S. B., Kaul, R. B., & Sutherland, D. M. (1991). New and corrected floristic records for Nebraska--4. *Transactions of the Nebraska Academy of Sciences*, 18, 141-150.

Rolfsmeier, S. B. (1995). Keys and distributional maps for Nebraska cyperaceae part 1: *Bulbostylis, cyperus, dulichium, eleocharis, eriophorum, fimbriostylis, fuirena, lipocarpha*, and *scirpus*. *Transactions of the Nebraska Academy of Sciences*, 22, 27-42.

Rosche, R. C., & Johnsgard, P. A. (1984). Birds of lake McConaughy and the North Platte river valley, Oshkosh to Keystone. *Nebraska Bird Review*, 52, 26-35.

Rose, K. E., Louda, S. M., & Rees, M. (2005). Demographic and evolutionary impacts of native and invasive insect herbivores on *Cirsium canescens*. *Ecology*, 86(2), 453-465.  
doi:10.1890/03-0697

Rowe, J. W. (1992). The occurrence of the sturgeon chub and introduced brook silverside in the Platte River of Nebraska. *Prairie Naturalist*, 24, 281-282.

Rowe, J. W. (1995). Hatchling size in the turtle *Chrysemys picta bellii* from western Nebraska: Relationships to egg and maternal body size. *Journal of Herpetology*, 29(1), 73-79.

Rowe, J. W. (1992). Observations of body size, growth, and reproduction in blanding's turtle (*Emydoidea blandingii*) from western Nebraska. *Canadian Journal of Zoology*, 70(9), 1690-1695. doi:10.1139/z92-235

Rowe, J. W. (1994). Reproductive variation and the egg size-clutch size trade-off within and among populations of painted turtles (*Chrysemys picta bellii*). *Oecologia*, 99(1-2), 35-44.  
doi:10.1007/BF00317081

Rowe, J. W. (1997). Growth rate, body size, sexual dimorphism and morphometric variation in four populations of painted turtles (*Chrysemys picta bellii*) from Nebraska. *American Midland Naturalist*, 138(1), 174-188.

Rowe, J. W., Holy, L., Ballinger, R. E., & Stanley-Samuelson, D. (1995). Lipid provisioning of turtle eggs and hatchlings: Total lipid, phospholipid, triacylglycerol and triacylglycerol fatty

acids. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 112(2), 323-330.

ROZEN JR, J.,G. (2000). Pupal descriptions of some cleptoparasitic bees (apidae), with a preliminary generic key to pupae of cleptoparasitic bees (apoidea). *American Museum Novitates*, 3289, 1-19. doi:10.1206/0003-0082(2000)289<0001:PDOSCB>2.0.CO;2

Rozen Jr, J.,G. (2013). Larval development and nesting biology of the adventive wood-nesting bee *Lithurgus (L.) chrysurus fonscolombe* (hymenoptera: Megachilidae: Lithurgini). *American Museum Novitates*, (3774), 20-40. doi:10.1206/3774.2

Rozen Jr, J. (1991). Evolution of cleptoparasitism in anthophorid bees as revealed by their mode of parasitism and first instars (hymenoptera: Apoidea). *American Museum Novitates*, 3029, 1-36.

Rozen, J. (2011). Descriptions of mature larvae of the bee tribe emphorini and its subtribes (hymenoptera, apidae, apinae). *Zookeys*, 148, 279-291. doi:10.3897/zookeys.148.1839

RUHNKE, T. R., & JAVONY JR, ,J. (1990). Life history differences between two species of *Gregarina* in *Tenebrio molitor* larvae. *The Journal of Parasitology*, 76(4), 519-522.

Russell, F. L., & Louda, S. M. (2004). Phenological synchrony affects interaction strength of an exotic weevil with platte thistle, a native host plant. *Oecologia*, 139(4), 525-534. doi:10.1007/s00442-004-1543-1

Russell, F. L., & Louda, S. M. (2005). Indirect interaction between two native thistles mediated by an invasive exotic floral herbivore. *Oecologia*, 146(3), 373-384. doi:10.1007/s00442-005-0204-3

Russell, F. L., Louda, S. M., Rand, T. A., & Kachman, S. D. (2007). Variation in herbivore-mediated indirect effects of an invasive plant on a native plant. *Ecology*, 88(2), 413-423. doi:10.1890/0012-9658(2007)88[413:VIHIEO]2.0.CO;2

Scharf, W. C. (2007). Woodland bird use of in-channel islands in the Central Platte River, Nebraska. *Prairie Naturalist*, 39(1), 15-28.

Scharf, W. C., Berigan, W. J., & Kren, J. (1993). Pine siskins in breeding condition along the North Platte River, Keith County, Nebraska, 1993. *Nebraska Bird Review*, 61(4), 144-145.

Scharf, W. C., & Kren, J. (1997). Summer diet of orchard orioles in southwestern Nebraska. *The Southwestern Naturalist*, 42(2), 127-131.

Schawang, J. E., & Janovy Jr, J. (2001). The response of *Gregarina niphandrodes* (apicomplexa: Eugregarinida: Septatina) to host starvation in *Tenebrio molitor* (coleoptera: Tenebrionidae) adults. *Journal of Parasitology*, 87(3), 600-605. doi:10.1645/0022-3395(2001)087[0600:TROGNA]2.0.CO;2

Schmidt-Rhaesa, A., Hanelt, B., & Reeves, W. K. (2003). Redescription and compilation of nearctic freshwater nematomorpha (gordiida), with the description of two new species. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 153(1), 77-117. doi:10.1635/0097-3157(2003)153[0077:RACONF]2.0.CO;2

Schotthoefer, A. M., Bolek, M. G., Cole, R. A., & Beasley, V. R. (2009). Parasites of the mink frog (*Rana septentrionalis*) from Minnesota, USA. *Comparative Parasitology*, 76(2), 240-246. doi:10.1654/4353.1

Scully, M. G. (2000). Mysterious creatures of a Nebraska lake. *Chronicle of Higher Education*, 47(5), B16-B16.

Shoop, W. L., & Janovy Jr, J.J. (1978). Adult cestodes from the coelomic cavity of the teiid lizard, *Cnemidophorus sexlineatus*. *Journal of Parasitology*, 64(3), 561-562.

Silcock, W. R., & Paseka, J. (2001). Summer field report, June and July 2001. *Nebraska Bird Review*, 69(3).

Smith, L. C., Raouf, S. A., Bomberger Brown, M., Wingfield, J. C., & Brown, C. R. (2005). Testosterone and group size in cliff swallows: Testing the "challenge hypothesis" in a colonial bird. *Hormones and Behavior*, 47(1), 76-82.

Snow, A. A., Pilson, D., Rieseberg, L. H., Paulsen, M. J., Pleskac, N., Reagon, M. R., . . . Selbo, S. M. (2003). A bt transgene reduces herbivory and enhances fecundity in wild sunflowers. *Ecological Applications*, 13(2), 279-286. doi:10.1890/1051-0761(2003)013[0279:ABTRHA]2.0.CO;2

Snyder, S. D., & Janovy Jr, J. (1994). Second intermediate host-specificity of *Haematoloechus complexus* and *Haematoloechus medioplexus* (digenea: Haematoloechidae). *The Journal of Parasitology*, 80(6), 1052-1055.

Snyder, S. D., & Janovy Jr, J. (1996). Behavioral basis of second intermediate host specificity among four species of *Haematoloechus* (digenea: Haematoloechidae). *The Journal of Parasitology*, 82(1), 94-99.

Snyder, S. R., & Friese, C. F. (2001). A survey of arbuscular mycorrhizal fungal root inoculum associated with harvester ant nests (*Pogonomyrmex occidentalis*) across the western United States. *Mycorrhiza*, 11(3), 163-165. doi:10.1007/s005720100119

Stanforth, L. M., Louda, S. M., & Bevill, R. L. (1997). Insect herbivory on juveniles of a threatened plant, *Cirsium pitcheri*, in relation to plant size, density and distribution. *Ecoscience*, 4(1), 57-66.

Stansbury, J., & Admiraal, D. M. (2004). Modeling to evaluate macrophyte induced impacts to dissolved oxygen in a tailwater reservoir. *Journal of the American Water Resources Association*, 40(6), 1483-1497.

Stanton-Geddes, J., & Anderson, C. G. (2011). Does a facultative mutualism limit species range expansion? *Oecologia*, 167(1), 149-155. doi:10.1007/s00442-011-1958-4

Steward, J. L., & Keeler, K. H. (1988). Are there trade-offs among antiherbivore defenses in *Ipomoea* (convolvulaceae)? *Oikos*, 53(1), 79-86.

Stoner, K. J. L., & Joern, A. (2004). Landscape vs. local habitat scale influences to insect communities from tallgrass prairie remnants. *Ecological Applications*, 14(5), 1306-1320. doi:10.1890/03-5112

Sutherland, D. M., & Kaul, R. B. (1986). Nebraska plant distribution. *Transactions of the Nebraska Academy of Sciences*, 14, 55-59.

Sutherland, D. M., & Rolfsmeier, S. B. (1989). An annotated list of the vascular plants of Keith County, Nebraska. *Transactions of the Nebraska Academy of Sciences*, 17, 83-101.

Sword, G. A., Joern, A., & Senior, L. B. (2005). Host plant-associated genetic differentiation in the snakeweed grasshopper, *Hesperotettix viridis* (orthoptera: Acrididae). *Molecular Ecology*, 14(7), 2197-2205. doi:10.1111/j.1365-294X.2005.02546.x

Sword, G. A., Senior, L. B., Gaskin, J. F., & Joern, A. (2007). Double trouble for grasshopper molecular systematics: Intra-individual heterogeneity of both mitochondrial 12S-valine-16S and nuclear internal transcribed spacer ribosomal DNA sequences in *Hesperotettix viridis* (orthoptera: Acrididae). *Systematic Entomology*, 32(3), 420-428. doi:10.1111/j.1365-3113.2007.00385.x

Tolley, G. S., & Hastings, V. S. (1960). Optimal water allocation: The North Platte River. *Quarterly Journal of Economics*, 74(2), 279-295.

Traxler, M. A., & Joern, A. (1999). Performance tradeoffs for two hosts within and between populations of the oligophagous grasshopper *Hesperotettix viridis* (Acrididae). [Host related differentiation among populations of the oligophagous grasshopper. (Acrididae) *Hesperotettix viridis*] *Oikos*, 87(2), 239-250.

Walker, M. H., Mackenzie, C., Bainbridge, S. P., & Orme, C. (1979). A study of the structure and gliding movement of *Gregarina garnhami*. *Journal of Eukaryotic Microbiology*, 26(4), 566-574. doi:10.1111/j.1550-7408.1979.tb04197.x

Watwood, S., Janovy Jr, J., Peterson, E., & Addison, M. A. (1997). *Gregarina triboliorum* (eugregarinida: Gregarinidae) n. sp. from *Tribolium confusum* and resolution of the confused taxonomic history of *Gregarina minuta* ishii, 1914. *Journal of Parasitology*, 83(3), 502-507.

Weller, S. G., Keeler, K. H., & Thomson, B. A. (2000). Clonal growth of *Lithospermum carolinense* (boraginaceae) in contrasting sand dune habitats. *American Journal of Botany*, 87(2), 237-242. doi:10.2307/2656911

Williams, J. A., & Nickol, B. B. (1989). Histological structure of the intestine and pyloric caeca of the green sunfish, *Lepomis cyanellus* rafinesque. *Journal of Fish Biology*, 35(3), 359-372. doi:10.1111/j.1095-8649.1989.tb02988.x

Winemiller, K. O., Pianka, E. R., Vitt, L. J., & Joern, A. (2001). Food web laws or niche theory? Six independent empirical tests. *The American Naturalist*, 158(2), 193-199.

Wise, M. R., Janovy Jr, J., & Wise, J. C. (2000). Host specificity in *Metamera sillasenorum*, n. sp., a gregarine parasite of the leech *Helobdella triserialis* with notes on transmission dynamics. *Journal of Parasitology*, 86(3), 602-606. doi: 10.1645/0022-3395(2000)086[0602:HSIMSN]2.0.CO;2

Wolesensky, W., Joern, A., & Logan, J. D. (2005). A model of digestion modulation in grasshoppers. *Ecological Modelling*, 188(2), 358-373.

Woodman, D. A. (1986). Microgeographic variation in *Rhinichthys cataractae* (cyprinidae) in western Nebraska. *Copeia*, 1986(1), 219-221.

Wythers, K., Wolfe, J., & Mole, S. (1993). Diet selection by larvae of the saltmarsh caterpillar in the Nebraska sandhills: Phenolics are not a problem. *Prairie Naturalist*, 25, 341-348.

Yang, Y., & Joern, A. (1994). Influence of diet quality, developmental stage, and temperature on food residence time in the grasshopper *Melanoplus differentialis*. *Physiological Zoology*, 67(3), 598-616.

Yang, Y., & Joern, A. J. (1994). Gut size changes in relation to variable food quality and body size in grasshoppers. *Ecology*, 8(1), 36-45.

Yang, Y., & Joern, A. (1994). Compensatory feeding in response to variable food quality by *Melanoplus differentialis*. *Physiological Entomology*, 19(1), 75-82. doi:10.1111/j.1365-3032.1994.tb01077.x